INSTRUCTIONS FOR USE ZORYVE[®] (zor-EEV) (roflumilast) topical foam, 0.3%

This Instructions for Use contains information on how to apply ZORYVE foam.

Read this Instructions for Use before you start using ZORYVE foam and each time you get a refill. There may be new information. This information does not take the place of talking with your healthcare provider about your medical condition or treatment. Use ZORYVE foam exactly as your healthcare provider tells you.

Important information you need to know before applying ZORYVE foam:

- **ZORYVE foam is for use on skin only (topical use).** ZORYVE foam is not for use in your eyes, mouth, or vagina.
- **ZORYVE foam is flammable.** Avoid fire, flame, and smoking during and right after you apply ZORYVE foam.


Before applying ZORYVE foam for the first time:



Gently pull back on the nozzle to break the plastic piece at the base.

Applying ZORYVE foam:

Apply a thin layer of ZORYVE foam 1 time a day to the affected areas of your body and scalp when they are not wet.



 Wash your hands after applying the medicine. If someone else applies ZORYVE foam for you, they should wash their hands after applying it.

Storing ZORYVE foam

- Store ZORYVE foam at room temperature between 68°F to 77°F (20°C to 25°C).
- ZORYVE foam is flammable. Keep away from heat and flame.
- The contents in ZORYVE foam are under pressure. **Do not** puncture or burn the can. **Do not** expose the can to heat or store at temperatures above 120°F (49°C).
- Do not freeze.
- Store the can upright.

Keep ZORYVE foam and all medicines out of the reach of children.

Marketed by:

Arcutis Biotherapeutics, Inc. Westlake Village, CA 91361

For more information on ZORYVE foam, call 1-844-692-6729 or visit http://www.zoryve.com.

This Instructions for Use has been approved by the U.S. Food and Drug Administration.

Revised: 5/2025

v2